

AD-A193 488 COMPARISONS BETWEEN SURFACE-ENHANCED RAMAN AND SURFACE
INFRARED SPECTROSC. (U) UTAH UNIV SALT LAKE CITY DEPT
UNCLASSIFIED OF CHEMISTRY S PONS ET AL. 30 JUL 86 TR-49
000044-02-K-0130

AD-A193 488 COMPARISONS BETWEEN SURFACE-ENHANCED RAMAN AND SURFACE
INFRARED SPECTROSC. (U) UTAH UNIV SALT LAKE CITY DEPT
UNCLASSIFIED OF CHEMISTRY S PONS ET AL. 30 JUL 86 TR-49
000044-02-K-0130


1/1


UNCLASSIFIED

UNCLASSIFIED

F/G 20/5

NL





MICROCOPY RESOLUTION TEST CHART
NBS 1963-A

AD-A193 488

OFFICE OF NAVAL RESEARCH

Contract N00014-83-K-0470-P00003

Task No. NR 359-718

TECHNICAL REPORT # 49

Comparisons Between Surface-Enhanced Raman and Surface
Infrared Spectroscopies for Strongly Perturbed
Adsorbates: Thiocyanate at Gold Electrodes

By

Stanley Pons
D. Corrigan
J. Foley
P. Gao
M. Weaver

Prepared for Publication in
Langmuir

University of Utah
Department of Chemistry
Salt Lake City, Utah 84112

July 30, 1986

Reproduction in whole or in part is permitted for
any purpose of the United States Government.

This document has been approved for public release
and sale; its distribution is unlimited.

DTIC
ELECTE
APR 14 1988
S D
H

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 49	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Comparisons Between Surface-Enhanced Raman and Surface Infrared Spectroscopies for Strongly Perturbed Adsorbates: Thiocyanate at Gold Electrodes		5. TYPE OF REPORT & PERIOD COVERED Technical Report # 49
7. AUTHOR(s) Stanley Pons, D. Corrigan, J. Foley, P. Gao, M. Weaver		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS University of Utah Department of Chemistry Salt Lake City, UT 84112		8. CONTRACT OR GRANT NUMBER(s) N00014-83-K-0470-P0003
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Chemistry Program - Chemistry Code 472 Arlington, Virginia 22217		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS Task No. NR 359-718
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE July 30, 1986
		13. NUMBER OF PAGES
		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) This document has been approved for public release and sale; its distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) IR spectroelectrochemistry, Surface Enhanced Raman, Thiocyanate		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The use of orthogonal collocation in the treatment of the problem of an array of ultramicroelectrodes is presented and compared to existing digital simulation techniques.		

LANGMUIR 1(1985) 616

COMPARISONS BETWEEN SURFACE-ENHANCED RAMAN AND SURFACE INFRARED SPECTROSCOPIES FOR STRONGLY PERTURBED ADSORBATES: THIOCYANATE AT GOLD ELECTRODES.

Dennis S. Corrigan,¹ John K. Foley,² Ping Gao,¹

Stanley Pons^{2*} and Michael J. Weaver^{1*}

Department of Chemistry, Purdue University
West Lafayette, IN 47907

and

Department of Chemistry, University of Utah
Salt Lake City, UT 84112

Abstract

The potential-dependent C-N stretching mode, ν_{CN} , for thiocyanate adsorbed at the gold-aqueous interface is examined by both surface-enhanced Raman spectroscopy (SERS) and subtractively normalized interfacial Fourier transform infrared spectroscopy (SNIFTIRS). Both SERS and SNIFTIRS exhibit a ν_{CN} band around 2120-2130 cm^{-1} over the potential range +500 to -500 mV vs. s.c.e., consistent with the presence of a predominantly S-bound adsorbate. The potential-dependent ν_{CN} frequencies were very similar (within ca. 5 cm^{-1}) for SNIFTIRS at smooth and electrochemically roughened gold, as well as for SNIFTIRS and SERS on the latter surface. This suggests that the SERS-active sites do not differ substantially from the preponderant sites sensed by SNIFTIRS. Some features of the SER spectra suggest that N- as well as S-bound adsorbed thiocyanate exists at far negative potentials.

There is currently rapid development occurring in the application of vibrational spectroscopic methods to the *in-situ* molecular characterization of electrochemical interfaces. Two such techniques are surface-enhanced Raman spectroscopy (SERS)³ and subtractively normalized interfacial Fourier transform infrared spectroscopy (SNIFTIRS).^{4,5} A virtue of the former approach is that the very large (ca. 10^5 - 10^7) surface enhancement of the Raman scattering, including unattached as well as surface-bound species,⁶ enable absolute vibrational spectra to readily be obtained at SERS-active surfaces, even for solutions containing high (≤ 0.1 M) bulk adsorbate concentrations. On the other hand, it is possible that adsorbate molecules at the particular surface sites that are primarily responsible for SERS exhibit atypical chemical and vibrational properties compared to those for the preponderant adsorbate molecules. Surface infrared spectroscopy does not suffer from this disadvantage since detectable spectra for adsorbed species can readily be obtained at smooth metal interfaces using SNIFTIRS or related difference spectral techniques, for which little or no surface enhancement of the infrared absorption appears to occur.⁴

This suggests that a valuable way of checking the applicability of SERS as a quantitative probe of surface structure is to compare surface Raman and infrared spectra obtained under identical conditions for adsorbates whose vibrational frequencies are known to be sensitive to the nature and type of the surface bonding involved. No such quantitative comparisons appear to have been reported previously.

In this communication we compare corresponding potential-dependent SERS and SNIFTIRS data gathered for the C-N stretching mode of thiocyanate adsorbed at gold electrodes, and summarize other SER spectral features for this system. The SNIFTIR spectra were obtained at both mechanically polished and electrochemically roughened surfaces, the latter also being

For	
&I	<input checked="" type="checkbox"/>
ed	<input type="checkbox"/>
tion	<input type="checkbox"/>
lon/	
ity Codes	
l and/or	
ocial	

A-1

COPY
INSPECTED

employed to acquire the SER spectra. Gold surfaces prepared by prior potential sweep oxidation-reduction cycles in chloride have recently been shown to provide remarkably stable and intense SER spectra for a miscellany of adsorbates in aqueous media.⁷ Thiocyanate is an interesting model adsorbate for several reasons, not the least of which are the large ($40\text{--}80\text{ cm}^{-1}$) increases in the C-N stretching frequencies accompanying metal-sulfur bonding, and the several spectral features that enable a distinction to be made between N- and S-coordination.⁸ Besides demonstrating close similarities in the potential-dependent C-N stretching frequencies for the corresponding surface Raman and infrared spectra, the present results provide an illustration of the merits of employing SERS and SNIFTIRS as complementary tools for gaining molecular structural information at electrode interfaces.

Experimental

Details of the SERS measurements are as given in ref. 9. Raman excitation was provided by a Spectra-Physics Model 165 Kr^+ laser operated at 647.1 nm, and the spectra gathered by using a SPEX Model 1403 double monochromator. SNIFTIRS measurements utilized an IBM Model 98-4A vacuum spectrometer, with the thin-layer electrochemical cell positioned in a small external compartment purged with nitrogen. Most details of the cell and the spectral measurements are given in ref. 5a.

The electrode used for the SERS measurements consisted of a 4 mm diameter gold disk sealed into a Teflon sheath of rotating disk construction (Pine Instruments). The SNIFTIRS electrode was a 7 mm gold disk sealed into Teflon. The electrodes were mechanically polished on a wheel with alumina down to 0.05 μm particle diameter and rinsed with water. The electrochemical roughening,

necessary for SERS, consisted of 20-30 potential sweeps between -300 mV and +1200 mV vs saturated calomel electrode (s.c.e.) in 0.1 M KCl. The surface remained shiny after this process, although gold electrodes displaying a pale brown hue and yielding even greater SERS intensities could be produced by holding the potential at the positive limit for 0.5-1 secs during each scan. Roughened as well as smooth electrodes having reflective properties were preferred for the SNIFTIRS measurements so to maximize the signal throughput. Differential capacitance measurements on these electrodes were undertaken as described in ref. 10. All electrode potentials are quoted versus the s.c.e., and all measurements were made at room temperature, $23 \pm 1^\circ\text{C}$.

Results and Discussion

Aqueous solutions containing ≥ 0.01 mM thiocyanate in 0.1 to 0.5 M Na_2SO_4 , NaClO_4 , or KCl supporting electrolytes yielded several SER spectral features at gold electrodes that indicate that NCS^- is specifically adsorbed over the entire polarizable potential range, +700 to -900 mV vs s.c.e. Representative SER spectra for 1 mM NaNCS in 0.5 M Na_2SO_4 at four electrode potentials are shown in Fig. 1. At the two most positive potentials, 500 and 100 mV (Fig. 1A,B), the spectra are similar to those seen previously for thiocyanate adsorbed at silver electrodes.^{11,12} Thus an intense and relatively broad ($\text{FWHM} \sim 30 \text{ cm}^{-1}$) C-N stretching band (ν_{CN}) is seen at $2110\text{-}2130 \text{ cm}^{-1}$, a weak band at about 700 cm^{-1} assigned to C-S stretching (ν_{CS}), a band around 450 cm^{-1} attributed to N-C-S bending (δ_{NCS}), and a broad feature around 240 cm^{-1} . The frequency of this last band is diagnostic of a metal surface-sulfur vibration ($\nu_{\text{Au-S}}$), SERS bands at similar frequencies being found at silver⁹ and gold¹³ electrodes for several N-bound metal thiocyanate complexes for which surface

attachment must occur via the sulfur atom. The low-frequency bands associated with surface-adsorbate vibrations for the supporting electrolyte anion, which occur at 185, 178, and 265 cm^{-1} for SO_4^{2-} , ClO_4^- , and Cl^- respectively,⁷ are completely removed upon addition of $\geq 0.1 \text{ mM}$ thiocyanate.

At more negative potentials ($\leq -100 \text{ mV}$, Fig. 1C,D), a new feature at 295 cm^{-1} appears and the 240 cm^{-1} band is reduced in intensity. This suggests that the S-bound thiocyanate is progressively being replaced by the N-bound form since these relative frequencies are consistent with that expected from the relative masses of the surface binding atoms.⁷

The behavior of the other vibrational modes are less conclusive in this regard. Thus the δ_{NCS} band remains at about 450-460 cm^{-1} throughout the complete potential range even though slightly higher frequencies would be anticipated if N-bound adsorbed thiocyanate is being formed.⁸ Although the ν_{CN} band decreases in intensity and shifts to progressively small frequencies in the range 2100-2130 cm^{-1} as the potential is made more negative, this frequency range is typical of that encountered for S- rather than N-bound thiocyanate, the latter most commonly being found around 2050-2080 cm^{-1} .⁸ In addition, a weak band around 700 cm^{-1} , also indicative of S-coordination, is obtained at positive potentials that weakens and eventually disappears at the most negative potentials.

Measurements of the differential double-layer capacitance against electrode potential ($C_{\text{dl}}-E$) also indicate the presence of substantial thiocyanate adsorption over the entire potential range 600 to -900 mV. This can readily be discerned from the addition of $\geq 0.1 \text{ mM}$ thiocyanate to 0.1 M KCl which results in loss of the broad $C_{\text{dl}}-E$ peak centered at 300 mV due to potential-dependent chloride adsorption. Smaller, roughly potential-independent capacitances, around 25 $\mu\text{F cm}^{-2}$, are obtained, which are indicative of high adsorbate coverages.¹⁰ The $C_{\text{dl}}-E$ curve shapes were essentially unaffected

by electrode roughening, although somewhat surprisingly this resulted in significantly (ca. 20%) *smaller* capacitance values.

The SNIFTIR spectra were obtained using -900 mV as the reference potential, where the extent of thiocyanate adsorption should be minimised. (Hydrogen evolution commences at more negative potentials.) Representative spectra obtained by stepping to several more positive potentials for 1 mM NCS^- in 0.5 M Na_2SO_4 at smooth gold are given in Fig. 2. Comparable results have also been obtained for similar conditions using electrochemically modulated infrared spectroscopy (EMIRS).^{4a,14} At the most negative potentials, ca. < -200 mV (Fig. 1A), a bipolar band occurs with positive- and negative-going peaks (features I and II) around 2105 and 2120 cm^{-1} , respectively. At more positive potentials the negative-going band broadens and shifts to higher frequencies, and a large positive-going band appears at around 2065 cm^{-1} (feature III, Figs. 1B-D).

Figure 3 shows a corresponding series of SNIFTIRS spectra for gold that was electrochemically roughened in the same manner as that used to produce the SER spectra in Fig. 1. These two sets of SNIFTIRS spectra are very similar; however, the broad negative-going band (feature II) on roughened gold is resolved into a pair of peaks at more positive potentials and a low frequency shoulder around 2040-2050 cm^{-1} appears on feature III (Fig. 3B,C).

Feature II is entirely consistent with the formation of additional S-bound thiocyanate as the potential becomes more positive. Figure 4 contains a plot of the SNIFTIRS peak frequencies, $\nu_{\text{CN}}^{\text{P}}$, of this band for smooth and roughened gold (closed triangles and circles, respectively) as a function of potential. (For the latter surface, $\nu_{\text{CN}}^{\text{P}}$ for the major, lower frequency, band is plotted.) Also included are the ν_{CN} frequencies obtained from the potential-dependent SER spectra on roughened gold (open symbols). Two sets of points are shown; the squares are the SERS

peak frequencies, $\nu_{\text{CN}}^{\text{P}}$, for the ν_{CN} band, whereas the circles are the "band center" (or average) frequencies, $\nu_{\text{CN}}^{\text{a}}$, obtained by bisecting the band area. [Small yet significant differences between $\nu_{\text{CN}}^{\text{P}}$ and $\nu_{\text{CN}}^{\text{a}}$ occur because the SERS bands are slightly asymmetric, $\nu_{\text{CN}}^{\text{P}}$ shifting from the high- to the low-frequency side of the band center as the potential becomes more negative (Fig. 1).]

Inspection of Fig. 4 shows that the SERS and SNIFTIRS ν_{CN} values are mostly within ca. 5 cm^{-1} of each other throughout the potential range 500 to -500 mV. The small differences are indeed comparable to the uncertainties in evaluating $\nu_{\text{CN}}^{\text{P}}$, especially for the infrared data. Admittedly, the potential dependence of $\nu_{\text{CN}}^{\text{P}}$ at roughened gold is significantly smaller for the SNIFTIRS than for the SERS data (Fig. 4). This apparent discrepancy is, however, probably due chiefly to the distortion of the negative SNIFTIRS ν_{CN} band (feature II) by its positive-going partner (feature I) since these two halves of the bipolar band overlap. Correction for this distortion will decrease $\nu_{\text{CN}}^{\text{P}}$ more at the more negative potentials where features II and I are least resolved. Nevertheless, the potential dependence of ν_{CN} appears to be relatively small; after allowing for these corrections one finds for both SERS and SNIFTIRS $d\nu_{\text{CN}}^{\text{P}}/dE \sim 12 \text{ cm}^{-1} \text{ V}^{-1}$ and for SERS $d\nu_{\text{CN}}^{\text{a}}/dE \sim 8 \text{ cm}^{-1} \text{ V}^{-1}$.

The values of $d\nu/dE$ are somewhat smaller than those found for several other structurally related systems using surface infrared or Raman measurements, such as NCS^- at silver,¹¹ CN^- at silver,¹⁵ or CO at Pt,¹⁶ although similar to that obtained for CN^- at Au using SERS, $15 \text{ cm}^{-1} \text{ V}^{-1}$.¹³ Some of the literature values may be influenced by potential-dependent variations in adsorbate coverage or structure. Given that an important component of these potential dependencies appears to be from an electric field (Stark) effect,¹⁷ the small $d\nu_{\text{CN}}/dE$ values for

NCS^- at gold may reflect the separation of the C-N group from the surface such that it experiences a smaller electrical field.

The large positive-going band at 2060 cm^{-1} (feature III) in the SNIFTIRS spectra has also been seen under similar conditions using EMIRS, and attributed chiefly to the presence of N-bound thiocyanate at negative potentials which reorients to the S-bound form at more positive potentials.^{4a,14} While this interpretation is qualitatively consistent with the low-frequency SERS data, as noted above there is an absence of SERS bands around 2060 cm^{-1} even at potentials as negative as -1000 mV . An alternative explanation is that feature III is associated chiefly with solution thiocyanate (for which $\nu_{\text{CN}} \approx 2060\text{ cm}^{-1}$) which is removed by adsorption as S-bound thiocyanate as the potential is altered from -900 mV to markedly more positive values. Although feature III is much more pronounced than the negative bands associated with S-bound thiocyanate (feature II, Figs. 2,3), this can be accounted for if the molar absorptivity of ν_{CN} for S-bound adsorbed thiocyanate, $\epsilon_{\text{NCS-}}$, is appropriately smaller than for uncoordinated thiocyanate, ϵ_{NCS} , as well as for the N-bound form, $\epsilon_{\text{SCN-}}$. At least for bulk-phase thiocyanate complexes, typically $\epsilon_{\text{NCS-}} \sim 0.2 \epsilon_{\text{NCS}} \sim 0.1 \epsilon_{\text{SCN-}}$;^{8,18} moreover, for $\text{Au}(\text{SCN})_4^-$, $\epsilon_{\text{NCS-}} \sim 0.1 \epsilon_{\text{NCS}}$.^{18a} Therefore the S-bound adsorption of a given amount of free thiocyanate from the thin solution layer is anticipated to yield a substantially (ca. tenfold) larger positive band at 2060 cm^{-1} than the accompanying negative SNIFTIRS band, in correspondence with the experimental data. The low-frequency shoulder on feature III at around $2040\text{--}2050\text{ cm}^{-1}$, seen most clearly with roughened gold (Fig. 3B,C) may well be due to reorientation of a small amount of N-bound thiocyanate present at -900 mV .

Admittedly, the assignment of the main feature III to loss of N-bound thiocyanate is in itself also in harmony with the data in Figs. 2 and 3. A

marked decrease in the intensity of feature III is, however, obtained when more dilute (0.1 mM NCS^-) solutions are employed and the electrode is pushed against the optical window so to minimise the thin-layer volume adjacent to the surface. This is consistent with the assignment of feature III to solution thiocyanate, but not with the adsorbed N-bound thiocyanate since the latter should be unaffected by the quantity of thiocyanate available for adsorption upon stepping to more positive potentials.

Taken together, then, the SERS and SNIFTIRS data both indicate the presence of S-bound thiocyanate at potentials less negative than ca. -400 mV , whereas some evidence suggests that the N-bound orientation may form at far negative potentials. The strong preference of the gold surface for the former thiocyanate orientation is expected both from chemical bonding¹⁹ and electrostatic arguments, the latter predicting S-bonding at positively charged surfaces since the sulfur atom carries most of the negative charge.⁸ Given that the potential of zero charge (p.z.c.) of polycrystalline gold is around $-50 \text{ mV vs. s.c.e.}$ in the absence of specific adsorption²⁰ and that a thiocyanate monolayer should shift this at least ca. 500 mV more negative,¹¹ extensive N-binding is expected to be restricted only to very negative potentials. It is also possible that both N- and S-binding occurs at more negative potentials with the thiocyanate lying flat on the surface; this geometry can account for the lack of a 2060 cm^{-1} peak in the SER spectra since such "bridging" thiocyanate usually yields a ν_{CN} band above 2100 cm^{-1} .⁸

Given the large ($60\text{--}70 \text{ cm}^{-1}$) increase of ν_{CN} upon S-bound adsorption of thiocyanate, the close (within 5 cm^{-1}) agreement in the frequencies of this species seen by the Raman and infrared probes, as well as the similarities of the latter at smooth and roughened electrodes, suggest

that the gold surface sites probed by SERS are chemically similar to the average or predominant sites that are presumably sensed by infrared spectroscopy. Close similarities have also been obtained between the SER and infrared spectra of cyanide on silver,^{4a,21} although parallel data sets using the same conditions for both techniques were apparently not obtained.

It will clearly be necessary, however, to examine systematically a number of systems under identical conditions in this manner before any sweeping conclusions can be drawn. Nevertheless, for substrates where Raman as well as infrared techniques can be employed, the strengths and limitations of the two methods are such that we anticipate that they will supply valuable complementary information.

Acknowledgments

This work is supported in part by grants from the Office of Naval Research (to SP and MJW) and from the Air Force Office of Scientific Research (MJW).

References

1. Purdue University
2. University of Utah
3. For recent reviews, see (a) Burke, R. L.; Lombardi, J. R.; Sanchez, L. A.; Adv. Chem. Ser. 1982, 201, 69; (b) Chang, R. K.; Furtak, T. E., Eds., "Surface-Enhanced Raman Scattering", Plenum, New York, 1982; (c) Chang, R. K.; Laube, B. L.; CRC Crit. Rev. Solid State Mater. Sci. 1984, 12, 1.
4. For recent reviews, see (a) Bewick, A.; Pons, S.; in "Advances in Infrared and Raman Spectroscopy", Hester, R. E., Clark, R., Eds, Heyden, London, 1984; (b) Bewick, A.; Pons, S.; in "Electroanalytical Chemistry", Bard, A. J., Ed., Marcel Dekker, New York, submitted.
5. (a) Pons, S.; Davidson, T.; Bewick, A.; J. Electroanal. Chem. 1984, 160, 63; (b) Pons, S.; J. Electroanal. Chem. 1983, 150, 495.
6. (a) Tadayyoni, M. A.; Farquharson, S.; Weaver, M. J.; J. Chem. Phys. 1984, 80, 1363; (b) Weaver, M. J.; Farquharson, S.; Tadayyoni, M. A.; J. Chem. Phys., in press.
7. Gao, P.; Patterson, M. L.; Tadayyoni, M. A.; Weaver, M. J.; Langmuir, 1985, 1, 173.
8. Bailey, R. A.; Kozak, S. L.; Michelson, T. W.; Mills, W. N.; Coord. Chem. Revs. 1971, 6, 407.
9. Tadayyoni, M. A.; Farquharson, S.; Li, T. T-T.; Weaver, M. J.; J. Phys. Chem. 1984, 88, 4701.
10. Hupp, J. T.; Larkin, D.; Weaver, M. J.; Surf. Sci. 1983, 125, 429.
11. Weaver, M. J.; Barz, F.; Gordon II, J. G.; Philpott, M. R.; Surf. Sci. 1983, 125, 409.
12. Wetzel, H.; Gerischer, H.; Pettinger, B.; Chem. Phys. Lett., 1981, 80, 159.
13. Gao, P., unpublished results.
14. Bewick, A., in preparation.
15. Kotz, R.; Yeager, E.; J. Electroanal. Chem. 1981, 123, 335.
16. Russell, J. W.; Severson, M.; Scanlon, K.; Overend, J.; Bewick, A.; J. Phys. Chem. 1983, 87, 293.
17. Lambert, D. K.; Solid State Comm., 1984, 51, 297; Korzeuiewski, K.; Pons, S.; Shirts, R. B.; J. Phys. Chem., in press.

18. (a) Bailey, R. A.; Michelson, T. W.; Mills, W. N.; J. Inorg. Nucl. Chem. 1971, 33, 3206; (b) Pecile, C.; Inorg. Chem. 1966, 5, 210.
19. Barclay, D. J.; Caja, J.; Croat. Chem. Acta, 1971, 43, 221.
20. Clavilier, J.; Huong, N.; Compt. Rend. 1971, 272, 1404; J. Electroanal. Chem. 1973, 41, 193.
21. Kanimatsu, K.; Seki, H.; Golden, W. G.; Chem. Phys. Lett., 1984, 108, 195.

Figure Captions

Figure 1

SER spectra for adsorbed thiocyanate at gold-aqueous interface at four electrode potentials as indicated (vs. s.c.e.). Solution contained 1 mM NaNCs in 0.5 M Na₂SO₄. Raman excitation was 50 mW spot focussed (ca. 2 mm diameter) on gold surface; spectral band-pass was 5 cm⁻¹. Typical peak intensity of ν_{CN} band was 3×10^3 counts sec⁻¹ versus background.

Figure 2


SNIFTIR spectra for thiocyanate at an electropolished gold-aqueous interface, using -900 mV vs s.c.e. as the reference potential, to four more positive potentials as indicated. Solution contained 1 mM NaNCs in 0.5 M Na₂SO₄. Spectra are an average of 1024 normalized scans at each potential.


Figure 3


SNIFTIR spectra for thiocyanate as in Figure 2, but using roughened gold under conditions used to generate the corresponding SER spectra in Figure 1.


Figure 4

The frequency of the C-N stretching mode for adsorbed thiocyanate at gold, ν_{CN} , obtained from SERS (open symbols) and SNIFTIRS (closed symbols) plotted against electrode potential, E. Circles are peak values of ν_{CN} obtained from SERS or SNIFTIRS data at roughened gold; closed triangles are corresponding SNIFTIRS data at smooth gold. Open squares are average values of ν_{CN} obtained from SERS (see text).









DL/413/83/01
GEN/413-2

TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 413 800 N. Quincy Street Arlington, Virginia 22217	2	Dr. David Young Code 334 NORDA NSTL, Mississippi 39529	1
Dr. Bernard Douda Naval Weapons Support Center Code 5042 Crane, Indiana 47522	1	Naval Weapons Center Attn: Dr. Ron Atkins Chemistry Division China Lake, California 93555	1
Commander, Naval Air Systems Command Attn: Code 310C (H. Rosenwasser) Washington, D.C. 20360	1	Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380	1
Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko Port Hueneme, California 93401	1	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, NC 27709	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1
DTNSRDC Attn: Dr. G. Bosmajian Applied Chemistry Division Annapolis, Maryland 21401	1	Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	1
Dr. William Tolles Superintendent Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375	1		

ABSTRACTS DISTRIBUTION LIST, 359/627

Dr. Paul Delahay
Department of Chemistry
New York University
New York, New York 10003

Dr. P. J. Hendra
Department of Chemistry
University of Southampton
Southampton SO9 5NH
United Kingdom

Dr. J. Driscoll
Lockheed Palo Alto Research
Laboratory
3251 Hanover Street
Palo Alto, California 94304

Dr. D. N. Bennion
Department of Chemical Engineering
Brigham Young University
Provo, Utah 84602

Dr. R. A. Marcus
Department of Chemistry
California Institute of Technology
Pasadena, California 91125

Dr. J. J. Auborn
Bell Laboratories
Murray Hill, New Jersey 07974

Dr. Joseph Singer, Code 302-1
NASA-Lewis
21000 Brookpark Road
Cleveland, Ohio 44135

Dr. P. P. Schmidt
Department of Chemistry
Oakland University
Rochester, Michigan 48063

Dr. Manfred Breiter
Institut für Technische Elektrochemie
Technischen Universität Wien
9 Getreidemarkt, 1160 Wien
AUSTRIA

Dr. E. Yeager
Department of Chemistry
Case Western Reserve University
Cleveland, Ohio 44106

Dr. C. E. Mueller
The Electrochemistry Branch
Naval Surface Weapons Center
White Oak Laboratory
Silver Spring, Maryland 20910

Dr. Sam Perone
Chemistry & Materials
Science Department
Lawrence Livermore National Laboratory
Livermore, California 94550

Dr. Royce W. Murray
Department of Chemistry
University of North Carolina
Chapel Hill, North Carolina 27514

Dr. B. Brummer
EIC Incorporated
111 Downey Street
Norwood, Massachusetts 02062

Dr. Adam Heller
Bell Laboratories
Murray Hill, New Jersey 07974

Dr. A. B. Ellis
Chemistry Department
University of Wisconsin
Madison, Wisconsin 53706

Library
Duracell, Inc.
Burlington, Massachusetts 01803

Electrochimica Corporation
20 Kelly Court
Menlo Park, California 94025-1418

ABSTRACTS DISTRIBUTION LIST, 359/627

Dr. M. Wrighton
Chemistry Department
Massachusetts Institute
of Technology
Cambridge, Massachusetts 02139

Dr. B. Stanley Pons
Department of Chemistry
University of Utah
Salt Lake City, Utah 84112

Donald E. Mains
Naval Weapons Support Center
Electrochemical Power Sources Division
Crane, Indiana 47522

S. Ruby
DOE (STOR)
Room 5E036 Forrestal Bldg., CE-14
Washington, D.C. 20595

Dr. A. J. Bard
Department of Chemistry
University of Texas
Austin, Texas 78712

Dr. Janet Osteryoung
Department of Chemistry
State University of New York
Buffalo, New York 14214

Dr. Donald W. Ernst
Naval Surface Weapons Center
Code R-33
White Oak Laboratory
Silver Spring, Maryland 20910

Mr. James R. Moden
Naval Underwater Systems Center
Code 3632
Newport, Rhode Island 02840

Dr. Bernard Spielvogel
U.S. Army Research Office
P.O. Box 12211
Research Triangle Park, NC 27709

Dr. Aaron Fletcher
Naval Weapons Center
Code 3852
China Lake, California 93555

Dr. M. M. Nicholson
Electronics Research Center
Rockwell International
3370 Miraloma Avenue
Anaheim, California

Dr. Michael J. Weaver
Department of Chemistry
Purdue University
West Lafayette, Indiana 47907

Dr. R. David Rauh
EIC Laboratories, Inc.
111 Downey Street
Norwood, Massachusetts 02062

Dr. Aaron Wold
Department of Chemistry
Brown University
Providence, Rhode Island 02192

Dr. Martin Fleischmann
Department of Chemistry
University of Southampton
Southampton SO9 5NH ENGLAND

Dr. R. A. Osteryoung
Department of Chemistry
State University of New York
Buffalo, New York 14214

Dr. John Wilkes
Air Force Office of Scientific
Research
Bolling AFB
Washington, D.C. 20332

Dr. R. Nowak
Naval Research Laboratory
Code 6171
Washington, D.C. 20375

Dr. D. F. Shriver
Department of Chemistry
Northwestern University
Evanston, Illinois 60201

ABSTRACTS DISTRIBUTION LIST, 359/627

Dr. Hector D. Abruna
Department of Chemistry
Cornell University
Ithaca, New York 14853

Dr. A. B. P. Lever
Chemistry Department
York University
Downsview, Ontario M3J1P3

Dr. Stanislaw Szpak
Naval Ocean Systems Center
Code 633, Bayside
San Diego, California 95152

Dr. Gregory Farrington
Department of Materials Science
and Engineering
University of Pennsylvania
Philadelphia, Pennsylvania 19104

M. L. Robertson
Manager, Electrochemical
and Power Sources Division
Naval Weapons Support Center
Crane, Indiana 47522

Dr. T. Marks
Department of Chemistry
Northwestern University
Evanston, Illinois 60201

Dr. Micha Tomkiewicz
Department of Physics
Brooklyn College
Brooklyn, New York 11210

Dr. Lesser Blum
Department of Physics
University of Puerto Rico
Rio Piedras, Puerto Rico 00931

Dr. Joseph Gordon, II
IBM Corporation
5600 Cottle Road
San Jose, California 95193

Dr. Nathan Lewis
Department of Chemistry
Stanford University
Stanford, California 94305

Dr. D. H. Whitmore
Department of Materials Science
Northwestern University
Evanston, Illinois 60201

Dr. Alan Bewick
Department of Chemistry
The University of Southampton
Southampton, SO9 5NH ENGLAND

Dr. E. Anderson
NAVSEA-56Z33 NC #4
2541 Jefferson Davis Highway
Arlington, Virginia 20362

Dr. Bruce Dunn
Department of Engineering &
Applied Science
University of California
Los Angeles, California 90024

Dr. Elton Cairns
Energy & Environment Division
Lawrence Berkeley Laboratory
University of California
Berkeley, California 94720

Dr. Richard Pollard
Department of Chemical Engineering
University of Houston
Houston, Texas 77004

Dr. M. Philpott
IBM Corporation
5600 Cottle Road
San Jose, California 95193

Dr. Donald Sandstrom
Boeing Aerospace Co.
P.O. Box 3999
Seattle, Washington 98124

Dr. Carl Kannewurf
Department of Electrical Engineering
and Computer Science
Northwestern University
Evanston, Illinois 60201

Dr. Joel Harris
Department of Chemistry
University of Utah
Salt Lake City, Utah 84112

ABSTRACTS DISTRIBUTION LIST, 359/627

Dr. Robert Somoano
Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California 91103

Dr. Johann A. Joebstl
USA Mobility Equipment R&D Command
DRDME-EC
Fort Belvoir, Virginia 22060

Dr. Judith H. Ambrus
NASA Headquarters
M.S. RTS-6
Washington, D.C. 20546

Dr. Albert R. Landgrebe
U.S. Department of Energy
M.S. 6B025 Forrestal Building
Washington, D.C. 20595

Dr. J. J. Brophy
Department of Physics
University of Utah
Salt Lake City, Utah 84112

Dr. Charles Martin
Department of Chemistry
Texas A&M University
College Station, Texas 77843

Dr. H. Tachikawa
Department of Chemistry
Jackson State University
Jackson, Mississippi 39217

Dr. Theodore Beck
Electrochemical Technology Corp.
3935 Leary Way N.W.
Seattle, Washington 98107

Dr. Farrell Lytle
Boeing Engineering and
Construction Engineers
P.O. Box 3707
Seattle, Washington 98124

Dr. Robert Gotscholl
U.S. Department of Energy
MS G-226
Washington, D.C. 20545

Dr. Edward Fletcher
Department of Mechanical Engineering
University of Minnesota
Minneapolis, Minnesota 55455

Dr. John Fontanella
Department of Physics
U.S. Naval Academy
Annapolis, Maryland 21402

Dr. Martha Greenblatt
Department of Chemistry
Rutgers University
New Brunswick, New Jersey 08903

Dr. John Wasson
Syntheco, Inc.
Rte 6 - Industrial Pike Road
Gastonia, North Carolina 28052

Dr. Walter Roth
Department of Physics
State University of New York
Albany, New York 12222

Dr. Anthony Sammells
Eltron Research Inc.
4260 Westbrook Drive, Suite 111
Aurora, Illinois 60505

Dr. C. A. Angell
Department of Chemistry
Purdue University
West Lafayette, Indiana 47907

Dr. Thomas Davis
Polymer Science and Standards
Division
National Bureau of Standards
Washington, D.C. 20234

Ms. Wendy Parkhurst
Naval Surface Weapons Center R-33
R-33
Silver Spring, Maryland 20910

DL/413/83/01
359/413-2

ABSTRACTS DISTRIBUTION LIST, 359/627

Dr. John Owen
Department of Chemistry and
Applied Chemistry
University of Salford
Salford M5 4WT ENGLAND

Dr. Boone Owens
Department of Chemical Engineering
and Materials Science
University of Minnesota
Minneapolis, Minnesota 55455

Dr. J. O. Thomas
University of Uppsala
Institute of Chemistry
Box 531
S-751 21 Uppsala, Sweden

Dr. O. Stafsudd
Department of Electrical Engineering
University of California
Los Angeles, California 90024

Dr. S. G. Greenbaum
Department of Physics
Hunter College of CUNY
New York, New York 10021

Dr. Menahem Anderman
W.R. Grace & Co.
Columbia, Maryland 20144

END

DATE

FILMED

DTIC

July 88